

# REMARKS ON OSMOSIS, QUANTUM MECHANICS, AND GRAVITY

ROBERT CARROLL  
UNIVERSITY OF ILLINOIS, URBANA, IL 61801

ABSTRACT. Some relations of the quantum potential to Weyl geometry are indicated with applications to the Friedmann equations for a toy quantum cosmology. Osmotic velocity and pressure are briefly discussed in terms of quantum mechanics and superfluids with connections to gravity.

## 1. REMARKS ON WEYL GEOMETRY

We begin with some features of Weyl geometry and the Dirac-Weyl theory following [54, 55, 56, 84] (cf. also [1, 2, 3, 4, 8, 12, 15, 16, 17, 19, 20, 23, 24, 25, 29, 77, 81, 85, 86, 87, 95]). Note that the paper is in part an expansion of some ideas in [24] and we correct some notational confusion from [17, 24]. There is a curious issue here of too much or too little, which is compounded by various, sometimes conflicting, notations and we feel that it is well advised to simply follow the extensive development of Israelit-Rosen which will keep the notation under control and benefit from insights therein. The background involves looking at a metric tensor  $g_{ab} = g_{ba}$  and a length connection vector  $w_\mu$ . If a vector is displaced by  $dx^\nu$  then

$$(1.1) \quad dB^\mu = -B^\sigma \hat{\Gamma}_{\sigma\nu}^\mu dx^\nu; \quad dB = B w_\nu dx^\nu$$

One can also write **(1A)**  $d(B^2) = 2B^2 w_\nu dx^\nu$  and this all requires that **(1B)**  $g_{\mu\sigma} \hat{\Gamma}_{\nu\sigma}^\sigma + g_{\nu\sigma} \hat{\Gamma}_{\mu\sigma}^\sigma = \partial_\sigma g_{\mu\nu} - 2g_{\mu\nu} w_\sigma$ . Then assuming that  $\hat{\Gamma}_{\mu\nu}^\sigma = \Gamma_{\mu\nu}^\sigma$  there results

$$(1.2) \quad \hat{\Gamma}_{\mu\nu}^\sigma = \Gamma_{\mu\nu}^\sigma + g_{\mu\nu} w^\sigma - \delta_\nu^\sigma w_\mu - \delta_\mu^\sigma w_\nu$$

where  $\Gamma_{\mu\nu}^\sigma$  denotes the standard Riemannian Christoffel symbol. The co-variant derivative for  $g_{ab}$  is written  $\nabla_\alpha$  or e.g.  $B_{;\nu}^\mu = \nabla_\nu B^\mu$  (and involves  $\Gamma_{\mu\nu}^\sigma$ ) whereas with the Weyl connection (1.2) one forms a co-covariant derivative of a vector

$$(1.3) \quad B_{;\nu}^\mu = \partial_\nu B^\mu + B^\sigma \hat{\Gamma}_{\sigma\nu}^\mu; \quad B_{\mu;\nu} = \partial_\nu B_\mu - B_\sigma \hat{\Gamma}_{\mu\nu}^\sigma$$

---

Date: April, 2011.  
email: rcarroll@math.uiuc.edu.

so that

$$(1.4) \quad B_{;\nu}^\mu = B_{;\nu}^\mu + B^\mu (g_{\sigma\nu} w^\mu - \delta_\nu^\mu w_\sigma - \delta_\sigma^\mu w_\nu)$$

Note also

$$(1.5) \quad g_{\mu\nu;\sigma} = 2g_{\mu\nu} w_\sigma; \quad g_{;\sigma}^{\mu\nu} = -2g^{\mu\nu} w_\sigma$$

Now under Weyl gauge transformations one has a length change (**1C**)  $B \rightarrow \hat{B} \sim \tilde{B} = e^{\lambda(x)} B$  with  $g_{\mu\nu} \rightarrow \tilde{g}_{\mu\nu} = e^{2\lambda} g_{\mu\nu}$  and  $g^{ab} \rightarrow e^{-2\lambda} g^{ab}$ . Writing  $e^{2\lambda(x)} = f(x)$  there is a calculation from [28] for lengths  $\ell \rightarrow \hat{\ell} = f^2 \ell$

$$(1.6) \quad 2\hat{\ell}d\hat{\ell} \sim f_\alpha dx^\alpha \ell^2 + 2f\ell d\ell \Rightarrow \frac{d\hat{\ell}}{\hat{\ell}} \sim \frac{1}{2} \partial_\alpha \log(f) dx^\alpha + w_\alpha dx^\alpha \Rightarrow \hat{w}_\alpha = \\ = w_\alpha + \frac{1}{2} \partial_\alpha \log(f) = w_\alpha + \partial_\alpha \lambda(x)$$

**REMARK 1.1.** In Weyl geometry consider now a scalar  $S$  of Weyl weight  $n$  defined via  $\tilde{S} = \exp(n\lambda)S$  (e.g.  $B = (g_{ab}B^aB^b)^{1/2} = \ell \rightarrow \hat{\ell} = e^\lambda \ell$ ) and  $\partial_\mu \tilde{S} = \exp(n\lambda)(\partial_\mu S + nS\partial_\mu \lambda)$ . This is not covariant with respect to Weyl gauge transformations (WGT) unless  $n = 0$  but one can define a gauge covariant derivative (**1D**)  $S_{||\mu} = \partial_\mu S - nS w_\mu$  with  $\tilde{S}_{||\mu} = \exp(n\lambda)S_{||\mu}$  (using (1.6)). For a gauge covariant derivative of a vector  $S^\mu$  of Weyl weight  $n$ , where  $\tilde{S}^\alpha = \exp(n\lambda)S^\alpha$ , there results (**1E**)  $S_{||\nu}^\mu = S_{;\nu}^\mu - nS^\mu w_\nu = \exp(n\lambda)S_{||\nu}^\mu$  (note  $\hat{\Gamma}_{\sigma\nu}^\mu - n\delta_\sigma^\mu w_\nu$  is actually a connection). ■

Dirac [28] (1973) introduced a  $\beta$  field of weight  $-1$  (as a Lagrange type multiplier) in order to have a satisfactory scalar tensor action of the form (cf. [54], Chap. 4)

$$(1.7) \quad I = \int \sqrt{-g} d^4x [-\beta^2 R + \sigma \beta^2 w^c w_c + (\sigma + 6) \partial_c \beta \partial^c \beta + 2\sigma \beta \partial_c \beta w^c + 2\Lambda \beta^4]$$

Note here that  $\sqrt{-g}$  is of weight 4 and  $R$  of weight -2; electromagnetic terms have been omitted (also  $\hbar = c = 1$ ). It is this factor  $\beta$  which was identified as a quantum mass  $\mathfrak{M} = m \exp(\mathfrak{Q})$  in [87] (and subsequently utilized in [15, 16, 17, 19, 20, 23, 24]). It's quantum significance was also noticed in [8] (cf. also [81]) in connection with conformal GR (GR stands for general relativity). Unfortunately a different notation was used in [8] and in [81] (0009169), namely (**1F**)  $\hat{g}_{ab} = \Omega^2(x)g_{ab}$  with  $\Omega^2 = \exp(-\psi)$  in [8] and  $\Omega^2 = \exp(\psi)$  in [81] (0009109) - a fact which I occasionally overlooked in subsequent work (cf. [24] for some errata). The latter paper [81] (0009169) contains some important material relating various actions in Einstein frame (EF) and Jordan frame (BD type equations). It was shown that an action

$$(1.8) \quad S_4 = \int d^4x \sqrt{-\hat{g}} e^{-\psi} \left[ \hat{R} - \left( \alpha - \frac{3}{2} \right) |\hat{\nabla} \psi|^2 + 16\pi e^{-\psi} L_M \right]$$

was the only one (of 4 considered) which was invariant under transformations of units (length, time, and mass) and this was then dubbed a string frame action (under the assumption that strings are ultimate and inevitable - cf. [44]). Note that  $\Omega^2 = \exp(\psi) = \phi$  implies

$$(1.9) \quad \hat{g}_{ab;c} = \partial_c \psi \hat{g}_{ab} \Rightarrow w_c = \frac{1}{2} \partial_c \psi$$

In particular conformal Riemannian geometry as in  $S_4$  is a WIST (Weyl integrable space time) as noted in [81] (0009169) and [73, 74].

Now  $S_4$  arises in our work via a conformal map  $g_{ab} \rightarrow \hat{g}_{ab} = \Omega^2 g_{ab}$  from an EF action (cf. [68]) **(1G)**  $S = \int \sqrt{-g} d^4x [R - \alpha |\nabla \psi|^2 + 16\pi L_M]$  in the form **(1H)**  $\hat{S} = \int \sqrt{-\hat{g}} d^4x [\hat{\phi} \hat{R} - (\omega/\hat{\phi}) |\hat{\nabla} \hat{\phi}|^2 + 16\pi \hat{\phi}^2 L_M]$  where we will write out the theory here in terms of  $\Omega^2 = \exp(\psi)$  following [19] (cf. also [15, 16] and note a few sign adjustments are needed in [17]). Recall first that  $\phi \sim \beta^2/m^2 = \mathfrak{M}^2/m^2$  and  $\hat{\phi} = \phi^{-1} = \exp(-\psi)$  and note that  $\sqrt{-g} = \hat{\phi}^2 \sqrt{-\hat{g}}$  (cf. [15, 16]) and

$$(1.10) \quad \sqrt{-\hat{g}} \hat{\phi} \hat{R} = \hat{\phi}^{-1} \sqrt{-\hat{g}} \hat{\phi}^2 \hat{R} = \hat{\phi}^{-1} \sqrt{-g} \hat{R} = \frac{\beta^2}{m^2} \sqrt{-g} \hat{R}$$

But via [28, 54] we can write (cf. also [17] for the pattern here)

$$(1.11) \quad \beta^2 \hat{R} = \beta^2 R - 6\beta^2 \nabla_\kappa w^\kappa + 6\beta^2 w^\kappa w_\kappa$$

and via  $-\beta^2 \nabla_{gk} w^\kappa = -\nabla_\kappa (\beta^2 w^\kappa) + 2\beta \partial_\kappa \beta w^\kappa$ , with a vanishing divergence term upon integration, the first integral in  $\hat{S}$  becomes

$$(1.12) \quad I_1 = \int \sqrt{-g} d^4x \left[ \frac{\beta^2}{m^2} R + 12\beta \partial_\kappa \beta w^\kappa + 6\beta^2 w^\kappa w_\kappa \right]$$

The second integral is ( $\gamma = \alpha - (3/2)$ )

$$(1.13) \quad I_2 = -\gamma \int \sqrt{-\hat{g}} d^4x \hat{\phi} \frac{|\hat{\nabla} \hat{\phi}|^2}{|\hat{\phi}|^2} = -\frac{4\gamma}{m^2} \int \sqrt{-g} d^4x |\hat{\nabla} \beta|^2$$

since  $\hat{\phi}_c = -\psi_c e^{-\psi}$ ,  $\hat{\nabla} \psi = -\hat{\nabla} \hat{\phi}/\hat{\phi}$ ,  $\hat{\phi} = \phi^{-1} = m^2 \beta^{-2}$ , and  $\hat{\phi}/\hat{\phi} = -2\hat{\nabla} \beta/\beta$  (along with (1.10)). Via **(1D)**  $\hat{\nabla}_c \beta = \partial_c \beta + w_c \beta$  for  $\Pi(\beta) = -1$  ( $\Pi$  denotes the Weyl weight) and from  $\beta^2 = m^2 \exp(\psi)$  one has  $\psi_c \exp(\psi) = (\beta^2/m^2) \psi_c \Rightarrow \psi_c = 2\beta_c/\beta$ . Hence via (1.9)  $w_c = \beta_c/\beta$  (correcting [17]) and consequently  $\hat{\nabla}_c \beta = 2\beta_c$ . In any event

$$(1.14) \quad \hat{S}_{GR} = \frac{1}{m^2} \int \sqrt{-g} d^4x [\beta^2 R + 12\beta \partial_c \beta w^c + 6\beta^2 w^c w_c - 4\gamma |\hat{\nabla} \beta|^2 + 16\pi m^2 L_M]$$

(as in [17], p. 59 (6.6) and p. 237 (3.32)).

**REMARK 1.2.** There was some sign confusion in [17] which we have adjusted above. Thus equations (6.7)-(6.8) on p. 59 and (3.33)-(3.34) on

p. 237 should be altered as indicated below. ■

First we write  $|\hat{\nabla}\beta|^2 = \beta^c\beta_c$  and from (1.9) and (1.14)

$$(1.15) \quad I_1 = \frac{1}{m^2} \int \sqrt{-g} d^4x [\beta^2 R + 18\partial_c\beta\partial^c\beta]; \quad I_2 = -16\gamma \int \sqrt{-g} d^4x \beta^c\beta_c$$

Consequently (recall  $\gamma = \alpha - (3/2)$ )

$$(1.16) \quad \hat{S} = \frac{1}{m^2} \int \sqrt{-g} d^4x [\beta^2 R + 16\pi m^2 L_M + 2(21 - 8\alpha)\beta^c\beta_c]$$

Then we rewrite (1.7) in the form

$$(1.17) \quad I = \int \sqrt{-g} d^4x [-\beta^2 R + 3(\sigma + 2)\beta_c\beta^c + 2\Lambda\beta^4]$$

Ignoring  $L_M$  and  $\Lambda$  for the moment we see that  $I = m^2\hat{S}$  provided that **(1I)**  $3\sigma = 8(3 - \alpha)$  and, a priori,  $\alpha$  or  $\sigma$  can be arbitrary (modulo e.g. possible cosmological restrictions on  $\alpha$ ).

**REMARK 1.3** The Brans-Dicke (BD) type theory of  $S_4$  (written also as  $\hat{S}_4$ ) was used for the Friedman equations in [24] for example where a relation  $\sigma \sim -4\alpha$  was used (based on some sign confusion). Hence one should convert the Weyl-Dirac action (1.1) of [24] to (1.7) and use the  $(\sigma, \alpha)$  relation **(1I)** in order to obtain the correct Friedmann equations for the toy quantum cosmology considered there (cf. Section 2). ■

## 2. REMARKS ON COSMOLOGY

In [24] the Friedmann equations were based on an  $S_4$  type action

$$(2.1) \quad \tilde{S} = \frac{1}{16\pi} \int \sqrt{-g} d^4x \left[ R\Phi - \omega \frac{|\nabla\Phi|^2}{\Phi} + L_M \right]$$

where **(2A)**  $L_M = -V(\Phi) + 16\pi\mathfrak{L}$  with  $\Lambda$  assumed to be suitably inserted in  $\mathfrak{L}_M$  (cf. [33, 38]). An FRW metric ( $c = 1$ )

$$(2.2) \quad ds^2 = dt^2 - a^2(t) \left[ \frac{dr^2}{1 - kr^2} + r^2 d\Sigma^2 \right]$$

is used and one has field equations as in [24]

$$(2.3) \quad G_{ab} = \frac{8\pi}{\Phi} T_{ab}^M + \frac{\omega}{\Phi^2} \left[ \nabla_a\Phi\nabla_b\Phi - \frac{1}{2}g_{ab}\nabla^c\Phi\nabla_c\Phi + \right. \\ \left. + \frac{1}{\Phi}(\nabla_a\nabla_b\Phi - g_{ab}\square\Phi) - \frac{V}{2\Phi}g_{ab} \right]$$

where **(2B)**  $T_{ab}^M = -(2/\sqrt{-g})(\delta/\delta g^{ab})(\sqrt{-g}\mathfrak{L}_M)$  (we ignore a possible shift  $\sqrt{-g} \rightarrow \sqrt{g}$ , etc. due to the signature in (2.2)). Variation of the action with respect to  $\Phi$  gives

$$(2.4) \quad \frac{2\omega}{\Phi} \square \Phi + R - \frac{\omega}{\Phi^2} \nabla^c \Phi \nabla_c \Phi - \frac{dV}{d\Phi} = 0$$

with trace

$$(2.5) \quad R = -\frac{8\pi T_M}{\Phi} + \frac{\omega}{\Phi^2} \nabla^c \Phi \nabla_c \Phi + \frac{3\square \Phi}{\Phi} + \frac{2V}{\Phi}$$

and using (2.5) to eliminate R from (2.4) yields

$$(2.6) \quad \square \Phi = \frac{1}{2\omega + 3} \left[ 8\pi T^M + \Phi \frac{dV}{d\Phi} - 2V \right]$$

Then using (2.2) one obtains **(2C)**  $\nabla^c \Phi \nabla_c \Phi = -(\dot{\Phi})^2$  and  $\square \Phi = -(\ddot{\Phi} + 3H\dot{\Phi}) = -(1/a^3)(d/dt)(a^3\dot{\Phi})$ . Assume now that **(2D)**  $T_{ab}^M = (P^M + \rho^M)u_a u_b + P^M g_{ab}$  and then the time dependent component of the BD field equations gives a constraint equation

$$(2.7) \quad H^2 = \frac{8\pi}{3\Phi} + \frac{\omega}{6} \left( \frac{\dot{\Phi}}{\Phi} \right)^2 - H \frac{\dot{\Phi}}{\Phi} - \frac{k}{a^2} + \frac{V}{6\Phi}$$

Then, using  $R = 6[\dot{H} + 2H^2 + (k/a^2)]$ , there results

$$(2.8) \quad \dot{H} + 2H^2 + \frac{k}{a^2} = -\frac{4\pi T^M}{3\Phi} - \frac{\omega}{6} \left( \frac{\dot{\Phi}}{\Phi} \right)^2 + \frac{1}{2} \frac{\square \Phi}{\Phi} + \frac{V}{3\Phi}$$

From (2.6), (2.7), and the trace equation **(2E)**  $T^M = 3P^M - \rho^M$  one has then

$$(2.9) \quad \begin{aligned} \dot{H} = & \frac{-8\pi}{(2\omega + 3)\Phi} [(\omega + 2)\rho^M + \omega P^M] - \frac{\omega}{2} \left( \frac{\dot{\Phi}}{\Phi} \right)^2 + \\ & + 2H \frac{\dot{\Phi}}{\Phi} + \frac{k}{a^2} + \frac{2}{2(2\omega + 3)\Phi} \left( \Phi \frac{dV}{d\Phi} - 2V \right) \end{aligned}$$

and (2.6) reduces to

$$(2.10) \quad \ddot{\Phi} + 3H\dot{\Phi} = \frac{1}{2\omega + 3} \left[ 8\pi(\rho^M - 3P^M) - \Phi \frac{dV}{d\Phi} + 2V \right]$$

In order to apply this to our model (1.7) we look at these Friedmann equations for  $\Phi = \hat{\phi}$  (with  $\mathfrak{Q} \sim \psi$ ) where  $(\beta^2/m^2) = \hat{\phi}^{-1} = \exp(\psi) = \exp(\mathfrak{Q})$ . Then  $\hat{\phi} = \exp(-\mathfrak{Q}) = \Phi$  and equations (2.3)-(2.10) can be written in terms of the quantum potential  $\mathfrak{Q}$  from **(2F)**  $\mathfrak{Q} = (\hbar^2/m^2)(\square|\Psi|/|\Psi|)$  with  $\Psi$  arising in the quantum process underlying (1.2) (cf. [2, 3, 4, 15, 16,

17, 20, 23, 24, 25, 85, 87, 95] and see also [6, 32]). We assume first that  $\mathfrak{L}_M = 0$  and then one can write  $(2\mathbf{G}) \dot{\Phi} = -\dot{\mathfrak{Q}}\Phi$ ,  $\ddot{\Phi} = (\dot{\mathfrak{Q}}^2 - \ddot{\mathfrak{Q}})\Phi$  and thence e.g.

$$(2.11) \quad (\dot{\mathfrak{Q}}^2 - \ddot{\mathfrak{Q}})\Phi - 3H\dot{\mathfrak{Q}}\Phi = \frac{1}{2\omega + 3} \left[ 8\pi(\rho^M - 3P^M) - \Phi \frac{dV}{d\Phi} + 2V \right]$$

Note that  $(2\mathbf{H}) \omega = \alpha - (3/2)$  and  $3\sigma = 8(3 - \alpha) \Rightarrow \sigma = 12 - (8/3)\omega$ . We see that the dynamics of  $\Phi$  is determined in part by  $V(dV/d\Phi) - 2V$  (which vanishes for  $V = c\Phi^2 = c\exp(-2\mathfrak{Q})$ ) and by  $\rho^M, P^M$ . Thus, in particular, it is possible to envision some cosmological behavior provided by a quantum background as in (1.7). We note also that Mannheim refers to intrinsically quantum mechanical gravity associated to general Weyl geometry with the Weyl tensor, etc. (cf. [5, 69]). We especially recommend Padmanabhan's paper [76] (1012.4476) and references there for a discussion of quantum mechanics and spacetime. Even more striking is the discussion in Grössing's book [46] (p. 123) which indicates (assuming an aether) how Einsteinian gravity can be considered a pure quantum phenomenon.

**REMARK 2.1.** It is interesting to note that via  $\hat{\phi} = \exp(-\mathfrak{Q})$  one has  $\mathfrak{Q} \sim \psi$  and hence by (1.9) the Weyl vector is  $w_c = (1/2)\partial_c\mathfrak{Q}$  implying a geometric role for  $\mathfrak{Q}$ . In particular this seems to show a direct connection of quantum mechanics with gravity. Of course this is not surprising since the geometry of (1.7) was generated via quantum mechanics but it suggests that Weyl geometry itself may have an implicit quantum connection via  $w_a = (1/2)\partial_a\mathfrak{Q}$  with  $\mathfrak{Q}$  a quantum potential. This seems to be a new observation (although connections of QM to the Weyl- Ricci curvature arise (cf. [2, 3, 15, 16, 20, 25, 85, 95])). ■

**REMARK 2.2.** In [87] there are also joint field equations involving the conformal factor  $\Omega^2$  (expressed via  $\phi$  or  $\phi^{-1}$ ) and for the quantum potential  $\mathfrak{Q}$  (expressed separately via a Lagrangian) with the goal of thereby deriving connections between  $\mathfrak{Q}$  and  $\phi$ . Also Bohmian aspects of the theory are emphasized. Some of this is also reviewed in [15, 16]. A main conclusion of this is that the quantum potential is a dynamical field and interactions between  $\Lambda$  and  $\mathfrak{Q}$  represent a connection between large and small scale structures. We note also from [73] that  $\Lambda$  plays an important role in generating mass for scalar fields and the understanding of  $\Lambda$  is considerably deepened and enhanced via work of Klinkhamer and Volovik (cf. [61, 62, 90, 91]) some of which is described briefly below. ■

**REMARK 2.3.** Relations between thermodynamics and gravity have been extensively studied following early work of Bekenstein and Hawking plus more recently Jacobson [57], and Padmanabhan [75, 76]. One can derive general Einstein field equations via thermodynamic principles and

gravity itself seems to be characterized via thermodynamics (see [17] for a brief sketch of some of this and there is much more information in works of Padmanabhan et al. The work of Verlinde [89] has triggered another explosion of interest in entropy and gravity and we mention here only a few articles, e.g. [9, 10, 13, 93] (related to the Friedmann equations) and [67, 70, 72, 88] (relations to quantum mechanics). ■

We would like to mention here that entropy in quantum mechanics is normally connected to momentum fluctuations and the quantum potential gives rise to an entropy functional  $\int P\mathfrak{Q} dx$  corresponding to Fisher information (here  $P = |\psi|^2$  where  $\psi$  is a wave function and  $\mathfrak{Q}$  represents a 3-dimensional quantum potential). Generally  $\mathfrak{Q}$  can be described e.g. via an osmotic velocity or a thermalization of this (cf. [15, 16, 17, 27, 34, 39, 45, 49, 50, 51] and Section 3). There is also a gravitational version of this related to the Wheeler-deWitt (WDW) framework in the form

$$(2.12) \quad \int \mathcal{D}h P\mathfrak{Q} = \int \mathcal{D}h \frac{\delta P^{1/2}}{\delta h_{ij}} G_{ijkl} \frac{\delta P^{1/2}}{\delta h_{kl}}$$

(cf. [16, 17, 21, 34, 51]) and see also [40, 92] for WDW). We note also the Perelman entropy functional

$$(2.13) \quad \mathfrak{F} = \int_M (R + |\nabla f|^2) e^{-f} dV$$

and corresponding Ricci flows are related to a so called Nash entropy **(2I)**  $S = \int u \log(u) dV$  where  $u = \exp(-f)$  and various aspects of quantum mechanics related to the Schrödinger equation and Weyl geometry arise (cf. [17, 18, 19, 43, 53, 59]).

### 3. SOME BACKGROUND

First it is interesting and perhaps enlightening to have some 3-D background here on the quantum potential. Thus, referring to [15, 16, 17, 22, 23, 26, 34, 35, 36, 37, 39, 45, 47, 48, 49, 50, 51, 71, 72, 82, 83, 88] we recall some results involving the QP relative to a 3-D SE. Various notations are used and we deal with a Schrödinger equation (SE) **(3A)**  $i\hbar\psi_t = \Delta\psi + V\psi$  with  $\psi = \text{Re} \exp(iS/\hbar)$  and

$$(3.1) \quad \partial_t S + \frac{1}{2m} |\nabla S|^2 + V + \mathfrak{Q} = 0; \quad \partial_t(R^2) + \frac{1}{m} \nabla(R^2 \nabla S) = 0$$

where the quantum potential (QP) is defined via

$$(3.2) \quad \mathfrak{Q} = -\frac{\hbar^2}{2m} \frac{\Delta R}{R} = -\frac{\hbar^2}{4m} \left[ \frac{1}{2} \left( \frac{\nabla P}{P} \right)^2 - \frac{\Delta P}{P} \right]$$

(note this is different from  $\mathfrak{Q}$  in e.g. (2F)). Here  $|\psi|^2 = R^2 = P$  and  $\rho = mP$ . Setting  $D = (\hbar/2m)$  one can write (3B)  $\mathbf{u} = -D(\nabla P/P)$  (osmotic velocity) which is expressed in terms of a “canonical” momentum fluctuation  $\delta p/m = \mathbf{u}$  (cf. [15, 16, 17] for references and more details). Now setting  $\mathbf{k}_u = -(1/2)\mathbf{u}$  one can write (3C)  $\mathfrak{Q} = (\hbar^2/2m)(|\mathbf{k}_u|^2 - \nabla \cdot \mathbf{k}_u) = (m/2)\mathbf{u}^2 - (\hbar/2)(\nabla \cdot \mathbf{u})$ . There is also a thermalization of the QP based on [45] (cf. also [17, 22]) whereby one refers to a basic particle energy  $E = \hbar\omega$  (where  $kT = \hbar\omega = 1/\alpha$  with  $k \sim k_B$  the Boltzman constant). One then defines a thermal function  $\mathcal{Q}$  where  $\Delta\mathcal{Q} = \mathcal{Q}(t) - \mathcal{Q}(0) < 0$  determines a heat dissipation with an osmotic velocity (3D)  $\mathbf{u} = -D(\nabla P/P) = (1/2\omega m)\nabla\mathcal{Q}$ . Hence and from (3.2), (3B), and (3D) there results

$$(3.3) \quad \int \mathfrak{Q} P d^3x = \frac{\hbar^2}{8m} \int \left( \frac{\nabla P}{P} \right)^2 P d^3x = \frac{m}{2} \int |\mathbf{u}|^2 P d^3x =$$

$$= \frac{1}{8\omega^2 m} \int |\nabla \mathcal{Q}|^2 P d^3x = \tilde{c}(t) \int (\nabla \mathcal{Q})^2 \exp^{-\alpha \mathcal{Q}} d^3x$$

(the last equation via  $\log(P) = -\alpha\mathcal{Q} + c(t) \Rightarrow P = \hat{c}(t)\exp(-\alpha\mathcal{Q})$  where  $\alpha = 1/\hbar\omega$  - cf. [17, 22, 45, 47, 48] for more on this). One should also note that (3E)  $\int P \mathfrak{Q} d^3x = (\hbar^2/8m) \int [(\nabla P)^2/P] d^3x = (\hbar^2/8m) FI$  where FI denotes Fisher information.

Now we note that an idea of osmotic pressure arises in the Klinkhamer-Volovik theory of superfluids (cf. [61, 62, 90, 91]) from which we extract here (see especially gr-qc 0711.3170, 0806.2805, 0907.4887, 0811.4347, and 1102.3152; hep-th 0907.4887 and 1101.1281; cond-mat 1004.0597; physics 0909.1044 and cf. also [30, 52, 79, 80]). This is a relativistic theory and the relativistic quantum vacuum is considered as a self sustained medium described by a variable  $\mathbf{q}$  which is a conserved quantity analogous to a particle density  $n$  in condensed matter theory but  $\mathbf{q}$  is the relativistic invariant quantity (see also Section 4). There is a Gibbs-Duhem relation  $\epsilon_{vac}(\mathbf{q}) - \mu\mathbf{q} = -P_{vac}$  where  $\epsilon_{vac}$  is the vacuum energy density and  $\mu$  is the vacuum chemical potential (thermodynamically conjugate to  $\mathbf{q}$ ). Dynamical equations for  $\mathbf{q}$  show that  $\Lambda = \epsilon_{vac} - \mu\mathbf{q} = -P_{vac}$  so the vacuum variable is automatically self-tuned to nullify in equilibrium any contribution from different quantum fields. In [91] (0909.1044) one considers a dilute solution of  ${}^3He$  quasiparticles in a superfluid vacuum  ${}^4He$  at  $T = 0$ . The negative contribution of the vacuum to the osmotic pressure of  ${}^3He$  is given by the same equation

$$(3.4) \quad P_{os} = P_{mat} + P_{vac} = P_{mat} - \frac{1}{2}\chi_{vac} \left[ \mathbf{q} \frac{\partial \epsilon_{mat}(n, \mathbf{q})}{\partial \mathbf{q}} \right]^2$$



where  $\chi_{vac}$  is the vacuum compressibility

$$(3.5) \quad \chi_{vac}^{-1} = \left[ \mathbf{q}^2 \frac{d^2 \epsilon_{vac}(\mathbf{q})}{d\mathbf{q}^2} \right]_{\mathbf{q}=\mathbf{q}_0} \geq 0$$

(where now  $\chi_{vac}$  corresponds to the compressibility of liquid  ${}^4\text{He}$ . This negative contribution to the osmotic pressure has been experimentally verified (according to 0909.1044).

**REMARK 3.1 .** Note that the formula in (3.3) involving the 3-D integral  $\int |\nabla \mathcal{Q}|^2 \exp(-\alpha \mathcal{Q}) d^3x$  reminds one of the  $I_2$  formula (1.13) from  $S_4$  where there is a 4-D integral involving  $\exp(-\psi) |\hat{\nabla} \psi|^2$ ; also the  $S_4$  or  $I_2$  represent “action” terms in the Weyl theory as does  $\int P \mathfrak{Q} d^3x$  in the 3-D theory (cf. [15, 16, 17]). However there is no immediate extrapolation here from 3-D to 4-D or vice-versa. In fact we know from remarks before (2.11) that  $\psi \sim \mathfrak{Q}$  in the 4-D theory where  $\mathfrak{Q}$  is given by (2F) and in the 3-D theory there is a formula (cf. [17, 45])

$$(3.6) \quad \mathfrak{Q} = -\frac{\hbar^2}{4k_B T m} \left[ \nabla^2 \mathcal{Q} - \frac{1}{D} \partial_t \mathcal{Q} \right]$$

We refer to Section 4 for some remarks on the KG equation and recall that there is related information via the Wheeler-deWitt (WDW) or Klein-Gordon (KG) equations in e.g. [6, 17, 21, 32, 34, 46, 51]. ■

**REMARK 3.2.** The role of information theory and entropy in QM has been clarified by e.g. Caticha [26] (see also in this connection [7, 71, 72, 88]). Osmotic velocity plays an important role in the diffusion theory of course and we mention also some recent work of Munkhammar [70] related to [89]. One can argue that sometimes QM is signaled simply by the entrance of  $\hbar$  into the theory. For example in the holographic force scenario of Verlinde [89] it seems to arise via the Unruh temperature relation to acceleration (3F)  $k_B T = (1/2\pi)(\hbar a/c)$ . Padmanabhan et al have written deeply and extensively about matters of entropy, entanglement, equipartition, horizons, and gravity as an emergent phenomenon (cf. [75, 76]). Entanglement entropy is also studied by J.W. Lee et al in [60, 63, 64] and one theme involves entropy and information erasure. An interesting Machian point of view is developed in [41, 42] and in [97] there is a microscopic approach to entropic gravity via the idea of coherence length. In [45] one defines a thermal term  $s$  via  $\nabla \mathcal{Q} = 2\omega \nabla(\delta s) = 2\omega m \mathbf{u} = -\omega \hbar (\nabla P/P)$ , providing the osmotic velocity, and in [26] this is combined with a relative entropy term  $\mathfrak{S}$ , based on “hidden” variables, for the “drift” velocity and a current velocity determined via  $\phi$  where  $\phi = \mathfrak{S} - \log(\rho^{1/2})$  with velocity connection  $v^a = b^a + u^a$

(see [26] for details). In [58] it is also shown that in entropic quantum dynamics gravitational potential and accelerating frames are informationally equivalent. ■

#### 4. SUPERFLUIDS

The superfluid universe of Klinkhamer and Volovik (cf. especially [91] (cond-mat 1004.0597) and [62] (0811.4347)) is a fascinating theme and we mention a few mathematical points here which were designed to give some structure to the superfluid world. We have a very imperfect and fragmentary idea of the physics here and only look at certain mathematical features which seem related to the material sketched above (and in [15, 16, 17] etc.). As an example of the arena for  $\mathfrak{q}$  theory one looks for  $\mathfrak{q}^{\mu\nu}$  which in a homogeneous vacuum has the form **(4A)**  $\mathfrak{q}^{\mu\nu} = \mathfrak{q}g^{\mu\nu}$  or **(4B)**  $\mathfrak{q}_{\mu\nu\alpha\beta} = \mathfrak{q}(g_{\alpha\mu}g_{\beta\nu} - g_{\alpha\nu}g_{\beta\mu})$  (generally here  $\hbar = c = k_B = 1$ ). Another example is **(4C)**  $\mathfrak{q}^{\mu\nu\alpha\beta} = \mathfrak{q}\epsilon^{\mu\nu\alpha\beta}$  where  $\epsilon^{\mu\nu\alpha\beta}$  is the fully antisymmetric Levi-Civita tensor (cf. [14]). More specifically one can consider a chiral condensate of gauge fields (e.g. a gluonic condensate in QCD). Assume e.g. that  $F_{\alpha\beta}$  represents a color magnetic field with **(4D)**  $\langle F_{\alpha\beta} \rangle = 0$  and  $\langle F_{\alpha\beta}F_{\mu\nu} \rangle = (\mathfrak{q}/24)\sqrt{-g}\epsilon_{\alpha\beta\mu\nu}$  where  $\mathfrak{q}$  is the anomaly driven topological condensate

$$(4.1) \quad \mathfrak{q} = \langle \tilde{F}^{\mu\nu}F_{\mu\nu} \rangle = \frac{1}{\sqrt{-g}}\epsilon^{\alpha\beta\mu\nu} \langle F_{\alpha\beta}F_{\mu\nu} \rangle$$

Then one chooses a vacuum action **(4E)**  $S_{\mathfrak{q}} = \int d^4x \sqrt{-g}\epsilon(\mathfrak{q})$  leading to a stress-energy tensor

$$(4.2) \quad T_{\mu\nu}^{\mathfrak{q}} = -\frac{2}{\sqrt{-g}}\frac{\delta S_{\mathfrak{q}}}{\delta g^{\mu\nu}} = \epsilon(\mathfrak{q})g_{\mu\nu} - 2\frac{\partial\epsilon}{\partial\mathfrak{q}}\frac{\partial\mathfrak{q}}{\partial g^{\mu\nu}} \Rightarrow \frac{\partial\mathfrak{q}}{\partial g^{\mu\nu}} = \frac{1}{2}g_{\mu\nu} \Rightarrow \\ \Rightarrow T_{\mu\nu}^{\mathfrak{q}} = g_{\mu\nu}\rho_{vac}(\mathfrak{q}); \quad \rho_{vac}(\mathfrak{q}) = \epsilon(\mathfrak{q}) - \mathfrak{q}\frac{\partial\epsilon}{\partial\mathfrak{q}}$$

In terms of the Einstein equations one has then **(4F)**  $T_{\mu\nu}^{\mathfrak{q}} = \Lambda g_{\mu\nu}$ ;  $\Lambda = \rho_{vac}(\mathfrak{q}) = \epsilon(\mathfrak{q}) - \mathfrak{q}\frac{\partial\epsilon}{\partial\mathfrak{q}}$ . Other possibilities are indicated in [62, 91] but fulfillment of (4.2) and **(4F)** is essential. Recall also from Section 3 that  $\epsilon_{vac} - \mu\mathfrak{q} = -P_{vac}$  where  $\mu$  is the chemical potential and one can argue that  $-P_{vac} \sim \Lambda \sim \rho_{vac}$ .

We recall now, from the discussion in Section 3, that given an osmotic velocity  $\mathbf{u}$  in some region  $\Omega \in \mathbf{R}^3$  there will be a local osmotic pressure

$$(4.3) \quad P_{loc} = \frac{1}{2}\rho\mathbf{u}^2 = \frac{m}{2}PD^2\frac{|\nabla P|^2}{P^2} = \frac{\hbar^2}{8m}\frac{|\nabla P|^2}{P}$$

so that **(4G)**  $P_{osm} = \int_{\Omega} P_u d^3x = \int \mathfrak{Q}P d^3x = (\hbar^2/8m)FI$ . In the superfluid universe mentioned above (with vacuum  ${}^4He$ ) one might suggest that

this could correspond to a vacuum osmotic pressure induced by a quantum generated “vacuum” based on Sections 1-2. In order to develop this one should examine the osmotic pressure idea in a 4-D context, in view of the expression (2F) for  $\Omega$ .

We sketch now from [46] (quant-ph 0201035) where the Nelsonian perspective is extended to a deBroglie-Bohm (BB) in the context of describing a KG framework (cf. also [65]). Perhaps somewhat prophetically it is mentioned in [46] (quant-ph 0201035) that the quantum vacuum has to play a major role in the evolution of the universe and one must take  $\Lambda$  seriously (which directive is being followed today, especially in the theory of superfluids). The introduction of an aether is appropriate in a relativistic (BB) theory and enters into the KG construction. We extract now from [46] (quant-ph 0201035) a few equations regarding osmotic velocity in hopes of implementing in some manner the superfluid theory. Thus, following [46], we write (4H)  $\psi(x, t) = R(x, t)\exp(-iS/\hbar)$ ;  $[\square + (m^2c^2/\hbar^2)]\psi = 0$  leading to  $(\square = (1/c^2)\partial_t^2 + \partial_x^2 + \partial_y^2 + \partial_z^2)$

$$(4.4) \quad \frac{\square R}{R} + \frac{1}{\hbar} \frac{\partial_\mu R}{R} \partial^\mu S + \frac{i}{\hbar} \frac{\partial^\mu R}{R} \partial_\mu S - \frac{1}{\hbar^2} \partial_\mu S \partial^\mu S + \frac{i}{\hbar} \square S + \frac{m^2 c^2}{\hbar^2} = 0$$

from which follows (for  $P = R^2$  and  $J_\mu = P \partial_\mu S$ )

$$(4.5) \quad 2 \frac{\partial^\mu R}{R} \partial_\mu S + \square S = 0; \quad \partial^\mu J_\mu = \partial^\mu P \partial_\mu S + P \square S = 0$$

along with (4I)  $\partial_\mu S \partial^\mu S = M^2 c^2 = m^2 c^2 + \hbar^2 (\square R/R)$ . Now to include vacuum energy one writes (4J)  $S^0 = E_{vac} t$  along with (4K)  $\Psi(x, t) = R(x, t)\exp[-i(S + S^0)/\hbar]$  for a solution of the KG equation. Then instead of (4.5) one involves a “hidden” probability current  $J_\mu^0 = P \partial_\mu S^0$  to get a conservation law for the total probability current (4L)  $\partial^\mu (J_\mu + J_\mu^0) = 0$  where (4M)  $\partial^\mu J_\mu = \partial^\mu P \partial_\mu S + P \square S = -\partial^\mu P \partial_\mu S^0 - P \square S^0 = -\partial^\mu J_\mu^0$ . Further  $S^0$  also modifies (4I) leading to a variable total mass

$$(4.6) \quad M^2 c^2 \rightarrow \left[ mc + \frac{E_{vac}}{c} \right]^2 + \hbar^2 \frac{\square R}{R}$$

Thus a (“hidden”) diffusion current  $J_\mu^0$  is introduced, due to the assumed stochastic aether dynamics and a corresponding all-pervading vacuum energy  $E_{vac}$ . One expects that timelike trajectories will ensue, i.e. the zero component  $J_{vac}^0 = P \partial_t (S + S^0) \geq 0$ .

Now in order to relate this causal KG theory with Nelson’s stochastic theory one defines (4N)  $S = Et - \mathbf{p}\mathbf{x} = \hbar p_\mu x^\mu$  such that  $\mathbf{v}^\mu = \partial^\mu S/M$ . Also an external potential  $V$  is assumed (which could include the vacuum energy  $E_{vac}$ ) with  $M$  given by (4I) as  $(\bullet) M = \{[m + (V/c^2)]^2 +$

$(\hbar^2/c^2)(\square R/R)\}^{1/2}$ . Thus the two real valued equations corresponding to the KG equation involves **(4O)**  $\partial^\mu J_\mu = \partial^\mu P \partial_\mu S + P \square S = 0$  and a Hamilton- Jacobi-Bohm equation

$$(4.7) \quad \partial_\mu S \partial^\mu S = M^2 \mathbf{v}_\mu \mathbf{v}^\mu = M^2 c^2 = \left[ mc + \frac{V}{c} \right]^2 + \mathfrak{Q}$$

where **(4P)**  $\mathfrak{Q} = \hbar^2(\square R/R)$ . This leads to an “osmotic velocity” **(4Q)**  $\mathbf{u}^\mu = D[\partial^\mu P/P]$  with  $D = \hbar/2m$  (cf. also [46] - book). One can then rewrite **(4O)** and (4.7) respectively as

$$(4.8) \quad \partial_\mu \mathbf{v}^\mu = -\frac{1}{D} \mathbf{u}_\mu \mathbf{v}^\mu - \frac{\partial_\mu M}{M} \mathbf{v}^\mu; \quad M^2 c^2 = \left[ mc + \frac{V}{c} \right]^2 + \mathfrak{Q}$$

where **(4R)**  $\mathfrak{Q} = m^2 c^2 + m \hbar \partial_\mu \mathbf{u}^\mu$  showing that the dynamics is essentially governed by the 4-gradient of the osmotic velocity. Also there is then a local osmotic pressure based on  $(1/2)M|\mathbf{u}|^2$  with M as in **(•)**. In the sense of [65] (but now in explicit dependence on the osmotic velocity) equations (4.8) are the two basic equations of relativistic stochastic mechanics (cf. [46] for more on this). Note that one can also rewrite (4.8-A) as

$$(4.9) \quad \partial_\mu \mathbf{v}^\mu = -\frac{1}{D}(\mathbf{u}_\mu + \mathbf{u}_\mu^M) \mathbf{v}^\mu; \quad \mathbf{u}_\mu^M = D \frac{\partial_\mu M}{M}$$

We refer to [46] for more equations and discussion and anticipate possible interesting connections of the analysis here to superfluid theory.

The obvious question now is to determine a possible relation between M and  $\mathfrak{M}$  where  $\mathfrak{M}$  is the quantum mass of Section 1 and M is the relativistic mass of [46]. We recall that the Weyl geometry involved  $\hat{g}_{ab} = \Omega^2(x)g_{ab}$  with **(4S)**  $\mathfrak{M}^2 = m^2 \exp(\mathfrak{Q})$  and **(4T)**  $\mathfrak{Q} = (\hbar^2/m^2)(\square|\Psi|/|\Psi|)$ . On the other hand there is no Weyl geometry explicit in Section 4 and one has from **(•)**, (4.7), (4.8)

$$(4.10) \quad M^2 c^2 = \left[ mc + \frac{V}{c} \right]^2 + \mathfrak{Q}; \quad \mathfrak{Q} = \hbar^2 \frac{\square R}{R}$$

Now we recall that when working directly with a KG equation it is natural to use an approximation **(4U)**  $\exp(\mathfrak{Q}) \sim 1 + \mathfrak{Q}$  (cf. [15, 16, 87]) and **(4S)** becomes then **(4V)**  $\mathfrak{M}^2 \sim m^2(1 + \mathfrak{Q})$ . Setting  $c = 1$  in (4.10) then one can compare

$$(4.11) \quad M^2 = (m + V)^2 + \hbar^2 \frac{\square R}{R} \quad \text{and} \quad \mathfrak{M}^2 = m^2 \left[ 1 + \frac{\hbar^2}{m^2} \frac{\square R}{R} \right]$$

Evidently  $\mathfrak{M}^2 \sim M^2$  with  $m \rightarrow m + V$  where V contains  $E_{vac}$ . If we take  $V = E_{vac}$  then M only differs from  $\mathfrak{M}$  by an  $E_{vac}$  term, i.e. for  $\tilde{\mathfrak{Q}} = \hbar^2 \square R/R$

$$(4.12) \quad \sqrt{M^2 - \tilde{\mathfrak{Q}}} - \sqrt{\mathfrak{M}^2 - \tilde{\mathfrak{Q}}} = E_{vac}$$

In a certain sense then the effect of relativity here is to display the influence of a quantum vacuum.

## REFERENCES

- [1] R. Adler, M. Bazin, and M. Schiffer, Introduction to general relativity, McGraw Hill, 1965
- [2] J. Audretsch, Phys. Rev. D, 24 (1981), 1470-1477; 27 (1983), 2872-2884
- [3] J. Audretsch, F. Gäehler, and N. Straumann, Comm. Math. Phys., 95 (1984), 41-51
- [4] J. Audretsch and C. Lämmerzahl, Class. Quant. Gravity, 5 (1988), 1285-1295
- [5] C. Bender and P. Mannheim, Phys. Rev. D, 78 (2008), 025022; Phys. Rev. Lett., 100 (2008), 110402; hep-th 0807.2607
- [6] G. Bertoldi, A. Faraggi, and M. Matone, Class. Quant. Grav., 17 (2000), 3965
- [7] D. Bohm and B. Hiley, Phys. Repts., 172 (1989), 93-122
- [8] R. Bonal, I. Quiros, and R. Cardenas, gr-qc 0010010
- [9] R. Cai and S. Kim, hep-th 05001055
- [10] R. Cai, L. Cao, and N. Ohta, hep-th 1001.3470 and 1002.1136
- [11] S. deCampo, R. Herrera, and P. Lebrana, gr-qc 0711.1559
- [12] V. Canuto, P. Adams, S. Hsieh, and E. Tsiang, Phys. Rev. D, 16 (1977), 1643-1663
- [13] Q. Cao, Y. Chen, and K. Shao, hep-th 1001.2597
- [14] M. Carmelli, Classical fields, World Scientific, 2001
- [15] R. Carroll, Fluctuations, information, gravity, and the quantum potential, Springer, 2006
- [16] R. Carroll, On the quantum potential, Arima Publ., 2007
- [17] R. Carroll, On the emergence theme of physics, World Scientific, 2010
- [18] R. Carroll, Progress in Physics, 4 (2007), 22-24; 1 (2008), 21-24 and 2 (2008), 89-90
- [19] R. Carroll, math-ph 0701077, 0703065, 0705.3921, 0712.3251; 0507027
- [20] R. Carroll, Foundations of Physics, 35 (2005), 131-154
- [21] R. Carroll, Teor. i Mat. Fiz., 152 (2007), 904-914
- [22] R. Carroll, math-ph 0807.1320 and 0807.4158
- [23] R. Carroll, Quantum Potential as Information: A mathematical survey, in: New trends in quantum information, Eds. Felloni, Singh, Licata, and Sakaji, Aracne Editrice, 2010, pp. 155-189
- [24] R. Carroll, math-ph 1007.4744; gr-qc 1010.1732
- [25] C. Castro, Found. Phys., 22 (1992), 569-615 and 37 (2007), 366-409; Found. Phys. Lett., 4 (1991), 81-99; Jour. Math. Phys., 31 (1990), 2626-2633
- [26] A. Caticha, quant-ph 0907.4335, 1005.2357, and 1011.0746
- [27] L. Crowell, Quantum fluctuations of spacetime, World Scientific, 2005
- [28] P. Dirac, Proc. Royal Soc. London, A 333 (1973), 403 and 338 (1974), 439
- [29] W. Drechsler, gr-qc 9901030
- [30] Z. Ezawa, Quantum Hall effects, World Scientific, 2008
- [31] F. Falciano, M. Novello, and J. Salim, Found. Phys., 40 (2010), 1885-1901
- [32] A. Faraggi and M. Matone, Inter. Jour. Mod. Phys. A, 15 (2000), 1869-2017
- [33] V. Faraoni, Cosmology in scalar-tensor theory, Kluwer, 2004; gr-qc 0906.1901 and 1005.2327
- [34] B. Frieden, Science from Fisher information, Springer, 2004; Physics from Fisher information, Cambridge Univ. Press, 1998
- [35] B. Frieden, A. Plastino, A. R. Plastino, and B. Soffer, Phys. Rev. E, 60 (1999), 48-53 and 66 (2002), 046128; Jour. Phys. A, 304 (2002), 73-78
- [36] B. Frieden and R. Gatenby, Exploratory data analysis using Fisher information, Springer, 2007

- [37] B. Frieden and B. Soffer, *Physica A*, 388 (2009), 1315-1330
- [38] Y. Fuji and K. Maeda, *The scalar-tensor theory of gravitation*, Cambridge Univ. Press, 2003
- [39] P. Garbaczewski, *Entropy*, 7 (2005), 253-299; *Jour. Stat. Phys.* 123 (2007), 315-355; quant-ph 0408192, 0612151, 0706.2481, and 0805.1536; cond-mat 0211362, 0604538, 0703147, 0703204, and 0811.3856
- [40] L. Glinka, gr-qc 0612079, 0707.3341, 0711.1380, 0712.1624, 0712.2769, 0801.4157, 0803.0533, and 0808.1533; physics 1102.5002 (book)
- [41] M. Gogberashvili, physics 0910.0169; gr-qc 0807.2439 and 1008.2544
- [42] M. Gogberashvili and I. Kanatchikov, gr-qc 0807.2439 and 1009.2266; physics 1012.5914
- [43] W. Graf, gr-qc 0602054
- [44] B. Greene, *The hidden reality*, Knopf, 2011
- [45] G. Grössing, *Entropy*, 12 (2010), 1975-2044; quant-ph 0201035, 0205048, 0311109, 0404030, 0410236, 0508079, 0711.4954, 0806.4462, and 0808.3539; *Phys. Lett. A*, 372 (2008), 4556-4562; *Found. Phys. Lett.*, 17 (2004), 343-362
- [46] G. Grössing, *Quantum cybernetics*, Springer, 2000; quant-ph 0201035 and 0205047
- [47] G. Grössing, S. Fussy, J. Mesa Pascasio, and H. Schwabl, quant-ph 1004.4596 and 1005.1058
- [48] G. Grössing and J. Mesa Pascasio, quant-ph 0812.3561
- [49] M. Hall, quant-ph 9802052, 9806013, 9903045, 9912055, 0103072, 0107149, 0302007; gr-qc 0408098; *Phys. Rev. A*, 55 (1997), 100-113 and 62 (2000), 012107, *Jour. Phys. A*, 37 (2004), 7799; *Gen. Relativ. Grav.*, 37 (2005), 1505-1515
- [50] M. Hall and M. Reginatto, *Jour. Phys. A* 35 (2002), 3289-3303; *Fortschrift Phys.*, 50 (2002), 646-651
- [51] M. Hall, K. Kumar, and M. Reginatto, *Jour. Phys. A*, 36 (2003), 9779-9794 (hep-th 0206235 and 0307259)
- [52] T. Harko, gr-qc 1101.3655
- [53] J. Isidro, J. Santander, and P. Fernandez de Cordoba, hep-th 0902.0143 and 0912.1535
- [54] M. Israelit, *The Weyl-Dirac theory and our universe*, Nova Sci. Publ., 1999
- [55] M. Israelit, *Found. Phys.*, 29 (1999), 1303; 32 (2002), 295 and 945; 28 (1998), 205; 35 (2005), 1725 and 1769; 37 (2007), 1628; gr-qc 9609035, 0710.3709, 0710.3910, 07103913, 07103923, and 0905.2482; astro-ph 1008.0767; *Inter. Jour. Mod. Phys. A*, 17 (2002), 4229
- [56] M. Israelit and N. Rosen, *Found. Phys.*, 13 (1983), 1023
- [57] T. Jacobson, *Phys. Rev. Lett.*, 75 (1995), 1260; hep-th 1001.4853; gr-qc 0711.3852 and 0801.1547
- [58] D. Johnson and A. Caticha, quant-ph 1010.1467
- [59] A. Kholodenko, gr-qc 0110064; hep-th 0303334 and 0701084
- [60] H. Kim, J. W. Lee, and J. Lee, 0709.3573
- [61] F. Klinkhamer, hep-th 1001.1939, 1006.2094, 1101.1281, and 1101.5370; gr-qc 0703009
- [62] F. Klinkhamer and G. Volovik, gr-qc 0711.3170, 0807.3896, 0811.4347, and 1102.3152; hep-th 0905.1919 and 0907.4887
- [63] J.W. Lee, hep-th 1003.1878, 1003.4464, 1005.2739, and 1011.1657
- [64] J.W. Lee, H. Kim and J. Lee, hep-th 0701199, 1001.5445 and 1002.4568
- [65] W. Lehr and J. Park, *Jour. Math. Phys.*, 18 (1977), 1235-1240
- [66] B. Liu, Y. Dai, X. Hu, and J. Deng, gr-qc 1012.2560

- [67] J. Makela, gr-qc 0506087, 0605098, 061078, 0805.3952, 0805.3955, 0810.4901, 1001.3808, and 1011.2052
- [68] G. Mangano and L. Sokolowski, gr-qc 9312008
- [69] P. Mannheim, hep-th 0707.2283, 0809.1200, 0909.0212, 0912.2635, 1005.5108, and 1101.2186; astro-ph 0505266 gr-qc 0001011 and 0703037
- [70] J. Munkhammar, hep-th 1003.1262 and 1006.0723; physics 1003.4981 and 1101.1417
- [71] M. Nagasawa, Schrodinger equations and diffusion theory, Birkhäuser, 1993; Stochastic processes in quantum physics, Birkhäuser, 2000
- [72] E. Nelson, Quantum fluctuations, Princeton Univ. Press, 1985; Dynamical theory of Brownian motion, PUP, 1967; Phys. Rev., 130 (1966), 1079-1085
- [73] M. Novello, astro-ph.CO, 1003.5126; physics 1004.3195 and 1008.2371
- [74] M. Novello, J. Salim, and F. Falciano, gr-qc 0901.3741; Found. Phys., 40 (2010), 1885-1901
- [75] T. Padmanabhan, Gravitation, Cambridge Univ. Press, 2010
- [76] T. Padmanabhan, gr-qc 0202078, 0209088, 0308070, 0311036, 0408051, 0412068, 0606061, 0910.0839, 0911.1403, 0911.5004, 0912.3165, 1001.3380, 1003.5662, 1007.5066, 1012.0119, and 1012.4476; hep-th 0205278
- [77] G. Papini, gr-qc 0111005 and 0304082
- [78] G. Papini and W. Wood, Phys. Lett. A, 202 (1995), 45-49
- [79] C. Pethick and H. Smith, Bose Einstein condensation in dilute gases, Cambridge Univ. Press, 2008
- [80] L. Pitaevskij and S. Stringari, Bose-Einstein condensates, Oxford Univ. Press, 2003
- [81] I. Quiros, gr-qc 9904004, 9905071, and 0004014; hep-th 0009169
- [82] M. Reginatto, Phys. Rev. A, 58 (1998), 1775; quant-ph 9909065; gr-qc 0501030
- [83] M. Reginatto and F. Lengyel, cond-mat 9910039
- [84] N. Rosen, Found. Phys., 12 (1982), 213
- [85] E. Santamato, Phys. Rev. D, 29 (1984), 216-222 and 32 (1985), 2615-2621; Jour. Math. Phys., 25 (1984), 2472-2480; Phys. Lett. A, 130 (1988), 199-202
- [86] E. Scholz, astro-ph 0403446 and 0805.2557; hep-th 1102.3478; gr-qc 0703102 and 0710.0269
- [87] F. and A. Shojai, gr-qc 0306099, 0404102, 0409053 and 001.3496; Inter. Jour. Mod. Phys. A, 15 (2000), 1859; Class. ,Quant. Grav., 21 (2004), 1-9
- [88] L. Smolin, hep-th 0605052; quant-ph 0609109; gr-qc 1001.3668; Class. Quant. Grav., 3 (1986), 347-359; Phys. Lett., 113A (1986), 408-412
- [89] E. Verlinde, hep-th 1001.0785
- [90] G. Volovik, The universe in a helium droplet, Oxford Univ. Press, 2003
- [91] G. Volovik, gr-qc 0004049, 0405012, 0406005, 0505104, and 0612134; cond-mat 0507454 and 1004.0597; physics 0909.1044; Found. Phys., 41 (2011), 516-528
- [92] C. Wang, gr-qc 0603062 and 0605124
- [93] S. Wei, Y. Liu, and Y. Wang, hep-th 1001.5238
- [94] S. Weinberg, Cosmology, Oxford Univ. Press, 2008
- [95] J. Wheeler, Phys. Rev. D, 41 (1990), 431-441 and 44 (1991), 1769-1773
- [96] W. Wood and G. Papini, Phys. Rev. D, 45 (1992), 3617-3627; Found. Phys. Lett., 6 (1993), 207-223; Phys. Lett. A, 202 (1995), 45-49; gr-qc 9612042
- [97] H. Xiong, hep-th 1012.5858, 1101.0525, and 1101.1270